



Brewing

New technologies

Edited by C. W. Bamforth

BARTH-HAAS GROUP

FlavorActiV™

KERRY
Kerry Bio-Science



WP

Brewing

Related titles:

Brewing: Science and practice

(ISBN-13: 978-1-85573-490-6; ISBN-10: 1-85573-490-7)

Brewing: Science and practice updates and revises the previous work of this distinguished team of authors, producing what is the standard work in its field. The book covers all stages of brewing from raw materials, including the chemistry of hops and the biology of yeasts, through individual processes such as mashing and wort separation. Key quality issues are discussed such as flavour and the chemical and physical properties of finished beers.

Cereal biotechnology

(ISBN-13: 978-1-85573-498-2; ISBN-10: 1-85573-498-2)

This major work explains the techniques involved and their enormous potential for food producers and consumers. It also explains how this new technology is regulated, the methods for assessing its potential risks, and the ways in which cereal biotechnology can add value.

Details of these books and a complete list of Woodhead titles can be obtained by:

- visiting our web site at www.woodheadpublishing.com
- contacting Customer Services (email: sales@woodhead-publishing.com; fax: +44 (0) 1223 893694; tel.: +44 (0) 1223 891358 ext. 30; address: Woodhead Publishing Limited, Abington Hall, Abington, Cambridge CB1 6AH, England)

Brewing

New technologies

Edited by
C. W. Bamforth

BARTH-HAAS GROUP

FlavorAct V™

KERRY
Kerry Bio-Science



CRC Press
Boca Raton Boston New York Washington, DC

WOODHEAD PUBLISHING LIMITED

Cambridge England

Published by Woodhead Publishing Limited, Abington Hall, Abington,
Cambridge CB1 6AH, England
www.woodheadpublishing.com

Published in North America by CRC Press LLC, 6000 Broken Sound Parkway, NW,
Suite 300, Boca Raton, FL 33487, USA

First published 2006, Woodhead Publishing Limited and CRC Press LLC
© 2006, Woodhead Publishing Limited
The authors have asserted their moral rights.

This book contains information obtained from authentic and highly regarded sources. Reprinted material is quoted with permission, and sources are indicated. Reasonable efforts have been made to publish reliable data and information, but the authors and the publishers cannot assume responsibility for the validity of all materials. Neither the authors nor the publishers, nor anyone else associated with this publication, shall be liable for any loss, damage or liability directly or indirectly caused or alleged to be caused by this book.

Neither this book nor any part may be reproduced or transmitted in any form or by any means, electronic or mechanical, including photocopying, microfilming and recording, or by any information storage or retrieval system, without permission in writing from Woodhead Publishing Limited.

The consent of Woodhead Publishing Limited does not extend to copying for general distribution, for promotion, for creating new works, or for resale. Specific permission must be obtained in writing from Woodhead Publishing Limited for such copying.

Trademark notice: Product or corporate names may be trademarks or registered trademarks, and are used only for identification and explanation, without intent to infringe.

British Library Cataloguing in Publication Data
A catalogue record for this book is available from the British Library.

Library of Congress Cataloging-in-Publication Data
A catalog record for this book is available from the Library of Congress.

Woodhead Publishing Limited ISBN-13: 978-1-84569-003-8 (book)
Woodhead Publishing Limited ISBN-10: 1-84569-003-6 (book)
Woodhead Publishing Limited ISBN-13: 978-1-84569-173-8 (e-book)
Woodhead Publishing Limited ISBN-10: 1-84569-173-3 (e-book)
CRC Press ISBN-13: 978-0-8493-9159-0
CRC Press ISBN-10: 0-8493-9159-8
CRC Press order number: WP9159

The publishers' policy is to use permanent paper from mills that operate a sustainable forestry policy, and which has been manufactured from pulp which is processed using acid-free and elementary chlorine-free practices. Furthermore, the publishers ensure that the text paper and cover board used have met acceptable environmental accreditation standards.

Project managed by Macfarlane Production Services, Dunstable, Bedfordshire, England
(e-mail: macfarl@aol.com)
Typeset by Godiva Publishing Services Ltd, Coventry, West Midlands, England
Printed by TJ International Limited, Padstow, Cornwall, England

Contents

<i>Contributor contact details</i>	xi
1 New brewing technologies: setting the scene	1
<i>C. W. Bamforth, University of California, USA</i>	
1.1 Introduction	1
1.2 The materials used in brewing	3
1.3 Brewing issues	8
1.4 Sources of further information	9
2 Providing cereals for brewing	10
<i>S. E. Heisel, American Malting Barley Association, USA</i>	
2.1 Introduction	10
2.2 Quality evaluations	11
2.3 Commercialization of new malting varieties	20
2.4 Future trends	23
2.5 References	25
3 Developments in the supply of adjunct materials for brewing .	30
<i>D. L. Goode, Kerry Bio-Science, The Netherlands and E. K. Arendt, University College Cork, Ireland</i>	
3.1 Introduction	30
3.2 Why use adjuncts?	30
3.3 The range of traditional adjuncts	32
3.4 Potential new adjunct sources	43
3.5 New developments to improve adjunct functionality	44
3.6 New beverages based on high adjunct levels	50

3.7	Future trends	57
3.8	Sources of further information	57
3.9	References	58
4	Malt and malt products	68
	<i>N. Davies, Muntons Plc, UK</i>	
4.1	Introduction – factors driving change	68
4.2	Malting barley development	70
4.3	Malt processing	76
4.4	Malted ingredients	92
4.5	Sensory evaluation of malt and malted ingredients	94
4.6	Future trends	97
4.7	Sources of further information	99
4.8	References	100
5	The breeding of hop	102
	<i>J. Henning, Oregon State University, USA</i>	
5.1	Introduction	102
5.2	Developments in hop breeding	104
5.3	Molecular techniques in hop cultivar development	106
5.4	Conclusion	118
5.5	Sources of further information	119
5.6	References	119
6	The processing of hops	123
	<i>C. Schönberger, Joh. Barth & Sohn GmbH & Co. KG, Germany</i>	
6.1	Introduction – the processing of hops	123
6.2	Hop pellets	125
6.3	Hop extracts	126
6.4	Isomerised hop products	132
6.5	Other hop products	138
6.6	Hop, hop products and relevant beer analyses	142
6.7	Future trends	145
6.8	Sources of further information	145
6.9	References	146
7	Yeast genetics in brewing: new insights and opportunities	149
	<i>D. E. Quain, red.ts Ltd, UK</i>	
7.1	Introduction	149
7.2	Fundamentals	151
7.3	The yeast genome	151
7.4	The rise and fall of genetic modification	154
7.5	Genetic instability – problem or opportunity?	157
7.6	Genetic differences between yeast species	159
7.7	Future trends	163

7.8	Sources of further information	163
7.9	Acknowledgements	163
7.10	References	163
8	Yeast supply and propagation in brewing	167
	<i>D. Quain, Coors Brewers Ltd, UK</i>	
8.1	Introduction	167
8.2	Storage and supply of yeast cultures	168
8.3	Propagation	171
8.4	Active dried yeasts	178
8.5	Future trends	179
8.6	Sources of further information	180
8.7	Acknowledgements	180
8.8	References	180
9	Water in brewing	183
	<i>M. Eumann, EUWA Water Treatment Plants, Germany</i>	
9.1	Introduction	183
9.2	Water for use in breweries	184
9.3	Treatment technologies	190
9.4	Recycling	205
9.5	Future trends	206
9.6	Sources of further information	206
9.7	Bibliography	207
10	The brewhouse	208
	<i>J. M. H. Andrews, Briggs of Burton Plc, UK</i>	
10.1	Introduction	208
10.2	Milling	209
10.3	Mashing devices	212
10.4	Mash conversion vessels	213
10.5	Mash separation	215
10.6	Wort boiling	221
10.7	Energy recovery systems	224
10.8	Hot wort clarification	225
10.9	Wort cooling	226
10.10	Cold wort clarification	226
10.11	Cold wort oxygenation	227
10.12	References	227
11	Fermentation of beer	228
	<i>C. Boulton, Coors Brewers Ltd, UK</i>	
11.1	Introduction	228
11.2	Current developments	229
11.3	Fermentation management	230

11.4	Yeast distribution in the fermenter	239
11.5	CO ₂ collection	241
11.6	Modelling fermentation	242
11.7	Continuous fermentation	244
11.8	High gravity brewing	246
11.9	Yeast physiology and fermentation performance	247
11.10	Future trends	249
11.11	Sources of further information	250
11.12	References	251
12	Accelerated processing of beer	254
	<i>I. Virkajärvi, VTT Technology, Finland</i>	
12.1	Introduction	254
12.2	Diacetyl reduction	255
12.3	New techniques	256
12.4	Warm stabilization	265
12.5	Cold stabilization	266
12.6	Two maturation philosophies	268
12.7	Sources of further information	269
12.8	References	270
13	Filtration and stabilisation of beer	275
	<i>G. Freeman, Brewing Research International, UK</i>	
13.1	Introduction	275
13.2	Current filtration practice	275
13.3	Possibilities in recycling of kieselguhr	278
13.4	Crossflow filtration	279
13.5	Single-pass membrane filtration	282
13.6	Novel filter aids	284
13.7	Current stabilisation processes	285
13.8	Novel stabilising systems	287
13.9	The effect of modern processing technologies on stability ...	288
13.10	Pasteurisation	289
13.11	Future trends	290
13.12	Acknowledgement	291
13.13	References	291
14	Packaging of beer	293
	<i>J. Browne, Technical Management Support Ltd, UK</i>	
14.1	Introduction	293
14.2	Trends in packaging formats	293
14.3	Developments in canning and bottling	298
14.4	Developments in handling kegs	303
14.5	Future trends	305
14.6	Sources of further information	307

14.7	Acknowledgements	307
14.8	References	307
15	Modern brewery sanitation	308
	<i>D. Loeffler, Loeffler Chemical Corporation, USA</i>	
15.1	Introduction	308
15.2	Sanitation terminology	309
15.3	Theoretical aspects of cleaning	311
15.4	The use of water in cleaning	313
15.5	Detergents/cleaners	314
15.6	Mechanical aspects of cleaning	319
15.7	Temperature	326
15.8	Time	327
15.9	Sanitizers/disinfectants	328
15.10	Future trends	333
15.11	Sources of further information	334
16	Waste handling in the brewing industry	335
	<i>R. Reed, Black & Veatch, UK</i>	
16.1	Introduction	335
16.2	Assessment of the character and strength of wastewaters	336
16.3	Sources and nature of wastewater	337
16.4	Extent of treatment	341
16.5	Outline of processes and technology	342
16.6	Treatment processes	346
16.7	Summary of the new legislation	353
16.8	Management approaches to water and waste minimisation ...	355
16.9	Future trends	356
16.10	References	357
17	Quality assurance in brewing	358
	<i>G. Jackson, Brewing Research International, UK</i>	
17.1	Introduction	358
17.2	Hazard Analysis Critical Control Points (HACCP)	358
17.3	Quality management systems and ISO9001:2000	363
17.4	Integrated management systems – the BRC Global Standard – Food	366
17.5	Feed Materials Assurance Scheme (FEMAS)	368
17.6	Future trends	371
17.7	References	371
18	Brewing control systems: chemical analytes	372
	<i>K. J. Siebert, Cornell University, USA</i>	
18.1	Introduction	372
18.2	Brewing analytical methods	373
18.3	Detection, separation and measurement techniques	373

18.4	Combining different techniques	379
18.5	What and why do we measure?	382
18.6	Where and how do we measure?	384
18.7	Impact of brewery operation scale	385
18.8	Changes over time	385
18.9	Traditional, emerging and future methods	386
18.10	New technologies	387
18.11	References	387
19	Brewing control systems: microbiological analysis	391
	<i>E. Storgårds, A. Haikara and R. Juvonen, VTT Technical Research Centre of Finland</i>	
19.1	Introduction	391
19.2	Classical microbiological methods	397
19.3	Optical techniques	400
19.4	Molecular methods for detection and identification	403
19.5	Indirect methods	407
19.6	Evaluation of yeast quality and quantity	409
19.7	Future trends	413
19.8	Sources of further information	415
19.9	References	416
20	Brewing control systems: sensory evaluation	427
	<i>W. J. Simpson, FlavorActiV Ltd, UK</i>	
20.1	Introduction	427
20.2	Brands	427
20.3	Tasters	431
20.4	Assessment methods	442
20.5	Assessment facilities	448
20.6	Data analysis tools	449
20.7	Data reporting and distribution tools	450
20.8	Future trends	450
20.9	Sources of further information	451
20.10	Acknowledgements	453
20.11	References	453
21	The future of brewing	461
	<i>C. W. Bamforth, University of California, USA</i>	
21.1	Choice	461
21.2	Impact of choice	462
21.3	Technical need drives research	462
21.4	Global influences	465
21.5	The nature of the brewing process in 2050	467
<i>Index</i>	468

Contributor contact details

(* = main contact)

Chapters 1 and 21

Dr C. W. Bamforth
Chair and Professor
Department of Food Science and
Technology
University of California
Davis, CA 95616
USA
E-mail: cwbamforth@ucdavis.edu

Chapter 2

Mr S. E. Heisel
American Malting Barley Association
740 N Plankinton Ave
Suite 830
Milwaukee, WI 53203-2403
USA

Chapter 3

Dr D. L. Goode*
Kerry Bio-Science BV
Veluwezoom 62
1327 AH Almere
The Netherlands
E-mail:
declan.goode@kerrybioscience.com

Dr E. K. Arendt
Department of Food and Nutritional
Sciences
University College Cork
Cork
Ireland
E-mail: e.arendt@ucc.ie

Chapter 4

Dr N. Davies
Technical Director
Muntons Plc
Cedars Maltings

Stowmarket
Suffolk IP14 2AG
UK
E-mail: nigel.davies@muntons.com

Chapter 5

Dr J. Henning
Research Hop Geneticist (plants)
USDA-ARS
3450 SW Campus Way
Corvallis, OR 97330
USA
E-mail: john.Henning@orst.edu

Chapter 6

Dr C. Schönberger
Joh. Barth & Sohn GmbH & Co. KG
Freiligrathstrasse 7/9
90482 Nuernberg
Germany
E-mail:
Christina.Schoenberger@johbarth.de

Chapters 7 and 8

Dr D. E. Quain
red.ts Ltd
9 Wheatfield Court
Willington
Derbyshire DE65 6PT
UK
E-mail: david.quain@red.ts.com

Chapter 9

M. Eumann
EUWA H.H. Eumann GmbH
Daimlerstrasse 2-10
D-71116 Gärtringen
Germany
E-mail: info@euwa.com

Chapter 10

Dr J. M. H. Andrews
Briggs of Burton Plc
Briggs House
Derby Street
Burton-on-Trent
Staffordshire DE14 2LH
UK
E-mail: jmha@briggsplc.co.uk

Chapter 11

Dr C. Boulton
Coors Brewers Ltd
Technical Centre
PO Box 12
Cross Street
Burton-on-Trent
Staffordshire DE14 1XH
UK
E-mail:
chris.boulton@coorsbrewers.com

Chapter 12

Dr I. Virkajärvi
Tuohimäki 3 B6
FIN-02130 Espoo
Finland
E-mail: ikka.virkajarvi@vtt.fi

Chapter 13

Mr G. Freeman
Brewing Research International
Coopers Hill Road
Nutfield
Redhill
Surrey RH1 4HY
UK

E-mail:
g.freeman@brewingresearch.co.uk

Chapter 14

Mr J. Browne
Technical Management Support Ltd
Claim Farm House
Manley Road
Kingswood
Frodsham
Cheshire WA6 6HT
UK

E-mail:
jeremy.browne@btinternet.com

Chapter 15

Mr D. Loeffler
Technical Director
Loeffler Chemical Corporation
5700 Bucknell Drive
Atlanta, GA 30336
USA

E-mail:
loeffler.dirk@loefflerchemical.com

Chapter 16

Dr R. Reed
Brackenbury House
Lodge Lane

Salfords
Redhill
Surrey RH1 5DH
UK
E-mail: reedr@bv.com

Chapter 17

Mr G. Jackson
Brewing Research International
Coopers Hill Road
Nutfield
Redhill
Surrey RH1 4HY
UK

E-mail:
g.jackson@brewingresearch.co.uk

Chapter 18

Dr K. J. Siebert
Food Science and Technology
Department
Cornell University
Geneva, NY 14456
USA

E-mail: kjs3@cornell.edu

Chapter 19

Dr E. Storgårds*, A. Haikara and
R. Juvonen
VTT Technical Research Centre of
Finland
PO Box 1000
FI-02044 VTT
Finland

E-mail: erna.storgards@vtt.fi

Chapter 20

Dr W. J. Simpson
FlavorActiV Limited
Sanderum House
Oakley Road
Chinnor
Oxfordshire OX39 4TW
UK

E-mail: bill.simpson@flavoractiv.com

Kerry Bio-Science, your brewing partner

Kerry Bio-Science is one of the market leaders in innovation and the application of bio- ingredients for brewing, distilling and beverage industries. Our range of brewing process aids and ingredients compliment the natural brewing process to produce beer of consistent quality. Our experienced team of experts help customers to create quality products that meet the rapidly changing demands of today's consumer market.

Kerry Bio-Science offers ingredients that can help you in every stage of the brewing process:

ENZYMES

Type of enzymes	: amylase, protease; glucanase for all brewing applications giving improved brewhouse performance (filtration and extract yield)
Functionality	
Range of products	: Bioglucanase, Bioamylase, Bioprotease, Promalt and Hitampase

FINING AGENTS

Functionality	: for wort and beer clarification
Range of products	: Whirlfloc and Bioline

FOAM STABILISERS

Functionality	: foam stabilisation
Range of products	: Biofoam

FLAVOUR STABILITY

Type of products	: anti-oxidant products
Range of products	: Blox

FERMENTATION AIDS

Functionality	: optimize fermentation performance
Range of products	: Yeastex and Fermcap



KERRY

Kerry Bio-Science

Brewing Technology Centre
Kerry Bio-Science Ireland
Kinagleary, Carrigaline
Co. Cork, Ireland
T +353 214 376400 / F +353 214 376480



1

New brewing technologies: setting the scene

C. W. Bamforth, University of California, USA

1.1 Introduction

The aim of this book is not primarily to tackle the science underpinning malting and brewing. Rather, the focus is on practical issues. In this chapter I will set the scene with some underpinning information, but those seeking the necessary basic scientific descriptions of everything from barley to beer should consult a text such as Bamforth (2006) or, at a more advanced level, Briggs *et al.* (2005). For immediate purposes Table 1.1 offers basic summaries of the processes involved in malting and brewing and the importance of each unit operation.

Table 1.1 The essentials of malting and brewing

Process stage	Description
<i>Selection of barley</i> ¹	Malting barleys (moisture content <12%) of relatively low total N content (e.g. less than 1.7% N), with high viability and with endosperm of mealy texture that hydrates readily and possesses cell walls that are readily degraded
<i>Storage of barley</i>	Sometimes important to allow barley to free up from innate dormancy
<i>Malting: steeping</i>	Staged addition of water at 14–18°C separated by air rests to raise moisture content to 43–46%. Allows the embryo to start synthesising hormones (notably gibberellins); allows the aleurone to become receptive to the hormone action that will trigger enzyme synthesis; allows the starchy endosperm to become receptive to digestion ('modification')

2 Brewing

Table 1.1 (continued)

Process stage	Description
<i>Malting: germination</i>	Controlled germination for 3–6 days at 16–20°C to degrade the endosperm cell walls and much of the protein. Enzymes synthesised by the aleurone migrate from proximal to distal end, digesting cell walls (β -glucanase family of enzymes, pentosanases) and protein (proteinase family). Causes softening of grain. Amylases also developed or activated, but limited in action on starch granules
<i>Malting: kilning</i>	Drying of malt at successively increasing temperatures (mainstream malts max. 105°C) to dry the grain (target <6% H ₂ O), whilst retaining much enzymic activity and developing colour and flavour through Maillard interactions between sugars and amino acids produced during modification
Malt: storage	2–4 weeks storage to avoid wort separation problems in the brewery
Milling and mashing	Generation of particles accessible to mashing water, mashing often starting at say 50°C (20 minutes) to allow remaining action of thermolabile β -glucanase, then passing through 65°C (e.g. for 1 h) for starch gelatinisation and action of amylase complex. Wort produced by conventional mashing has 20–25% starch left behind as non-fermentable digestion products called dextrans. Wort separated from grains
Boiling	Wort boiled with hops or hop preparations, typically for 1 hour. Isomerisation of bitter acids (main part of resin fraction) to increase their solubility and bitterness. Volatilisation of aroma components, including hop oils unless hops added late in the boil. Clarification stage follows to remove ‘hot break’ and residual hop material. Wort cooled and air or oxygen added
Fermentation	Ales typically fermented warmer (15–25°C) and therefore faster than lagers (6–15°C). Time range 3–14 days. Fermentation at a targeted rate of specific gravity drop and to a target ‘attenuation’. Also diacetyl and pentanedione, which afford butterscotch/honey aromas, must be removed by prolonged contact of yeast with ‘green beer’
Maturation, stabilisation and packaging	Minimum regime is –1°C, usually for 2–3 days. Some hold longer. Insolubilisation and settling of proteins and polyphenols. Filtration (kieselguhr or perlite-based). Removal of residual haze precursors by polyvinylpyrrolidone (polyphenols) and/or silica hydrogels or tannic acid or papain (proteins). Removal of any microbial contamination by pasteurisation or filtration. Adjustment of CO ₂ content, then fill vessels

¹ *Process stages in italics occur in the maltings, prior to the brewery.*

Source: Bamforth, C.W. (2004) Opportunities for newer technologies in the oldest biotechnology, brewing. *Applied Biotechnology, Food Science and Policy*, 1, 213–222.

1.2 The materials used in brewing

The larger proportion of the world's beers is produced from *malted barley*. It has been this way for perhaps 8000 years. Barley has been retained as the primary cereal of choice, not least because it retains its husk on threshing and this traditionally form the filter bed through which wort is collected in the brew house. As we shall see in Chapter 10, the mash filter does not depend on the husk in this way, which might in future open up possibilities for huskless barleys for brewing. However, these handle in the maltings much as does *wheat*, with a tendency towards stickiness. Wheat is the second most employed cereal in brewing, notably for the production of weissbiers and weizenbiers in Germany.

Barley is malted prior to use, in order that the enzymes that degrade starch to fermentable sugars are synthesised. Additionally there is a synthesis of the enzymes that degrade the cell walls and much of the protein in the starchy endosperm, thereby softening the grain and making it more millable. There is an unavoidable development of embryo tissue (rootlets and acrospire), and the maltster seeks to balance the extent of this with the need for adequate 'modification' of the endosperm.

Although there are beers that are produced only from malted barley e.g. those produced under the terms of the five-centuries-old Bavarian 'purity' law, the *Reinheitsgebot*, many brewers employ various adjuncts for reasons of quality (different colours, better foams, interesting flavours) or cost. In fact there may not be quite the cost savings anticipated from use of an ostensibly cheaper starch source, because the brewer may have to employ more expensive processing procedures if problems are to be avoided. The major cost components in brewing are illustrated in Fig. 1.1. It will be apparent that the costs of packaging, processing, taxation and marketing vastly exceed those of raw materials. It is only when there are political reasons or taxation reasons – e.g. Happoshu, see Chapter 3) – that there are major justifications to be made on a cost basis for the use of adjuncts.

Of particularly low relative cost to the brewer are the *hops*, yet these make a huge contribution to product quality and stability: apart from the bitterness from the resins and aroma from the oils, the resins also afford anti-microbial properties and foam stabilisation but also comprise precursors of staling and of the skunky aromas that develop in beer exposed to light. Despite the low cost of hops relative to that of the beer, there is a plethora of hop preparations available to the brewer, including those that specifically protect against this light sensitivity (Chapter 6).

The bulk of most beers comprises *water*, hence the scrutiny which brewers devote to this product. Vastly more water is needed to make a pint of beer than actually finds its way into the beer, perhaps five times more for a well-run brewery and 20 times more for a badly operated facility.

Most brewers maintain their own *yeast* strains. As the alcohol concentration of most beers does not become too high during fermentation, the yeast that multiplies in fermentation remains healthy and suitable for re-pitching into



Fig. 1.1 A generalised breakdown of major costs associated with the brewing of beer.

subsequent fermentations. Some smaller brewers have done this in seeming perpetuity; however, the latter-day 'gold standard' is to re-pitch with newly propagated yeast every fourth or fifth fermentation. There is some interest in the use of dried yeast preparations of the type used *inter alia* in baking and wine making (Chapter 8).

As they will not be addressed elsewhere in this book, let us now consider miscellaneous process ingredients that may be employed in brewing. Not included in the discussion are the salts that may be added to adjust the ionic balance of the grist, e.g. calcium salts to increase the hardness ('Burtonisation'), and the zinc chloride or zinc sulphate that many brewers add to encourage yeast activity.

1.2.1 Exogenous enzymes

Various 'brewing enzymes' have been marketed over the years with the intent of satisfying various demands of the brewer (Table 1.2). The majority of these enzymes are not pure, insofar as they contain a range of enzymic activities. During the growth of the organism, different enzymes will be successively released into the medium in the order in which they are needed. So the precise balance of enzymes that is packaged and made available to the brewer will very much depend on where the 'cut' was taken. Nowadays most of the brewing enzymes are made using organisms that are genetically modified so as to optimise the level and type of enzymes in the broth.

1.2.2 Isinglass

One of the great beer genres (*viz.* the English cask ale) emerged on the backbone of a 'natural' clarification process rooted in a protein preparation called isinglass. Isinglass is a very pure form of collagen obtained from the dried swim bladders (some call them 'maws') of certain warm-water fish, amongst them the catfish, jewfish, threadfish and croaker. These fish are primarily caught for food use and the functional property of the maw represents added value.

Table 1.2 Exogenous enzymes used in brewing

Enzyme	Stage of use	Function
β -Glucanase ¹	Mashing	Eliminate glucans that cause wort separation, filtration and clarity problems
Pentosanase ¹	Mashing	Support glucanases in digestion of cell wall polymers from barley and wheat
Proteinase ¹	Mashing	Ensure generation of sufficient yeast-assimilable amino acids
α -Amylase ¹	Mashing	Starch digestion
Glucoamylase	Mashing or fermentation	Enhance starch digestion, to the extent of allowing conversion to totally fermentable sugars of value in the production of light and low carbohydrate beers
Pullulanase	Mashing	Promote digestion of branched-chain dextrins
Acetolactate decarboxylase	Fermentation	Accelerate elimination of vicinal diketones in beer maturation
Papain	Beer in storage	Eliminate haze-forming polypeptides
Prolyl endopeptidase	Beer in storage	Selectively remove haze-forming polypeptides; of potential value in producing beer for coeliacs

¹ Of especial significance if malt replaced by high levels of grain adjunct, e.g. unmalted barley.

The bladders are removed, washed and dried. At the smallest scale in a fishing village the maws are sun-dried, but modern fish processing plants use commercial dryers. Dried maws are ground up, washed and sterilised before being 'cut' by weak acids such as sulphurous acid to disrupt the structure of the collagen molecules so as to generate the correct balance and orientation of positively and negatively charged sites that are responsible for its functionality.

Worldwide, few brewers use isinglass, despite the fact that it is very effective in settling the contents of conditioning tanks, thereby minimising the loading of solids onto the filter without the expense of a centrifuge.

Various other fining agents are sometimes used in the brewing industry. These may be classified as kettle (or copper) finings, which are added during the boil, and auxiliary finings, which are used alongside isinglass to aid clarification of beer. The best-known kettle fining agent is the negatively charged carrageenan (Irish Moss), derived from seaweed, and the most frequently used auxiliary finings are silicates and alginates, the former derived from sand and the latter from seaweed. Again, both are negatively charged and complement the action of the positively charged isinglass.

1.2.3 Filter aids

There are fundamentally two types of filter aid: kieselguhr or perlite. Kieselguhr, or 'diatomaceous earth', comprises silica-based shells of ancient unicellular aquatic microscopic plants called diatoms. Its heat resistance means that it can

be used as an insulator, but its abrasiveness means that it has also formed a component of toothpaste and metal polishes. Apart from being widely used as a filter aid to clarify syrups as well as alcoholic beverages, it is used as a filling material in paper, paints, ceramics, soap and detergents. Alfred Nobel found that it is a great absorbant of nitro-glycerine in the manufacture of dynamite.

Huge beds of kieselguhr, between 40 and 50 feet (12–15 metres) deep, are found in Virginia, but also in parts of Germany and in Aberdeenshire in Scotland. The microscopic appearance from different localities differs considerably. The deposits contain varying amounts of organic matter together with sand, clay, and iron oxide, and the raw material is first incinerated (calcined) to destroy organic matter. The successive process stages in rendering bags of kieselguhr in the form needed by the brewer are mining, crushing, drying, calcining, cooling, air classification and packaging.

The interests of purveyors of perlite are looked after by the Perlite Institute (go to <http://www.perlite.org/>). Perlite is a naturally occurring siliceous rock that, when heated, expands from four to 20 times its original volume. When heated to above 871°C, it pops like popcorn to produce many small bubbles, so perlite is very light and white.

There are many uses for perlite. Its insulating properties and lightness render it valuable as an insulator in masonry and cryogenic vessels. It is used as an aggregate in cement and plasters and for under-floor insulation, chimney linings, paint texturing, gypsum boards, ceiling tiles, and roof insulation boards. Perlite is used as a component of soil-less growing mixes, allowing aeration and moisture retention. It is also used as a carrier for fertiliser, herbicides and pesticides and for pelletising seed.

Apart from clarifying beer, perlite is also used for cleaning up pharmaceuticals, chemicals and water. Like kieselguhr, it can also be used as an abrasive.

1.2.4 Stabilisers

Silica hydrogels and xerogels have their origins in a pure form of sand. This is first converted into a soluble form by the action of alkali. Thereafter there is a controlled aggregation of sodium silicate from a sol form under acidic conditions. The washed aggregated particles are processed by techniques including micronisation, drying, milling and classifying to yield the desired balance of particle sizes and pore sizes, the range of which is taken advantage of by brewers (and others) to remove colloidal particles of various types from their products.

Polyvinylpyrrolidone (PVPP) is produced by a technique called 'popcorn polymerisation'. The monomer vinylpyrrolidone is heated with strong caustic and then cooled, in which phase the polymerisation takes place. Subsequently there are slurring, filtering, hydrolysis (using phosphoric acid), washing, re-slurring, and drying stages, in which any residual monomer and water are removed.

Alternative stabilisers are now in the market, including a PVP–silica gel composite.

1.2.5 Foam stabiliser

Propylene glycol alginate (PGA) is rather more relevant as a foam *protectant* than as a foam stabiliser. It helps prevent lipid and detergent damage at point of sale, without offering absolute protection. Alginates are polysaccharides extracted by alkali from brown seaweed. They have been known since their discovery by an English chemist in 1881. The seaweed is harvested on the coasts of North America, Scotland, Ireland, Norway, France, Japan, China, Korea, Chile, South Africa and Australia. Of these China easily generates the most, though the UK and the USA are the biggest processors, with one company dominating, at 70% of the market.

Insoluble alginate from the raw plant is solubilised by ion exchange, then filtered. The alginic acid obtained is partially dehydrated and then esterified by reaction with gaseous propylene oxide under pressure at 45–60°C. After washing with alcohol, the product is dried and milled.

Table 1.3 Issues of quality in brewing

Issue	Description
Foam	Most beers are expected to have an appealing, stable foam. However, increasing tendency for consumers to drink directly from the container. Impacting factors include amphipathic polypeptide survival from the grist, presence of bitter acids (reduced side-chain acids give more stable foams), levels of CO ₂ and N ₂ , nucleation devices, absence of lipids and detergents.
Clarity	Most beers are bright (free from haze or sediment) – but not all. Insoluble materials can have diverse origins, including proteins cross-linking with oxidised polyphenols, residual grain polysaccharides, oxalate, micro-organisms
Colour	Derives from Maillard reaction in malt kilning and adjunct roasting and from polyphenol oxidation in the brew house
Absence of gushing	Principal cause is <i>Fusarium</i> infection of grain, with release of hydrophobin. But can also be cause by oxidised resins in hop bitterness preparations and insolubles in beer, e.g. oxalate
Package	Major determinant of impression customer gets of product – e.g. absence of scuffing on bottles, making returnable glass less appealing than one-trip glass
Flavour	Impacted by all the above. Pronounced marketing pressure to have beer in any coloured glass other than brown, but flint and green glass permit light passage that causes skunking of beer. Flavour impacted by all raw materials, demanding control of material selection and processing. Flavour instability the major unsolved problem facing brewers, in particular development of papery flavour. Stringent control of oxygen a major factor, but not alone. Cold shipping and stockholding of beer expensive but effective

Note: enhanced analytical capabilities (e.g. sensors) will permit increased control, with impacts on cost and quality.

sample content of Brewing: New Technologies (Woodhead Publishing Series in Food Science, Technology and Nutrition)

- [Lacanian Ethics and the Assumption of Subjectivity pdf, azw \(kindle\)](#)
- [read online Cocaine Wars](#)
- **[download The Art of The Last of Us](#)**
- [read Ivar's Seafood Cookbook: The O-fish-al Guide to Cooking the Northwest Catch](#)

- <http://www.gateaerospaceforum.com/?library/Beer-O-Clock--An-Insider-s-Guide-to-History--Craft--and-Culture.pdf>
- <http://aircon.servicessingaporecompany.com/?lib/The-Souls-of-Black-Folk--Oxford-World-s-Classics-.pdf>
- <http://honareavalmusic.com/?books/The-Art-of-The-Last-of-Us.pdf>
- <http://reseauplatoparis.com/library/The-Original-Curse--Did-the-Cubs-Throw-the-1918-World-Series-to-Babe-Ruth-s-Red-Sox-and-Incite-the-Black-Sox-Sc>