



Brewing

New technologies

Edited by C. W. Bamforth

BARTH-HAAS GROUP

FlavorActiV™

KERRY
Kerry Bio-Science



WP

Brewing

Related titles:

Brewing: Science and practice

(ISBN-13: 978-1-85573-490-6; ISBN-10: 1-85573-490-7)

Brewing: Science and practice updates and revises the previous work of this distinguished team of authors, producing what is the standard work in its field. The book covers all stages of brewing from raw materials, including the chemistry of hops and the biology of yeasts, through individual processes such as mashing and wort separation. Key quality issues are discussed such as flavour and the chemical and physical properties of finished beers.

Cereal biotechnology

(ISBN-13: 978-1-85573-498-2; ISBN-10: 1-85573-498-2)

This major work explains the techniques involved and their enormous potential for food producers and consumers. It also explains how this new technology is regulated, the methods for assessing its potential risks, and the ways in which cereal biotechnology can add value.

Details of these books and a complete list of Woodhead titles can be obtained by:

- visiting our web site at www.woodheadpublishing.com
- contacting Customer Services (email: sales@woodhead-publishing.com; fax: +44 (0) 1223 893694; tel.: +44 (0) 1223 891358 ext. 30; address: Woodhead Publishing Limited, Abington Hall, Abington, Cambridge CB1 6AH, England)

Brewing

New technologies

Edited by
C. W. Bamforth

BARTH-HAAS GROUP

FlavorAct V™

KERRY
Kerry Bio-Science



CRC Press
Boca Raton Boston New York Washington, DC

WOODHEAD PUBLISHING LIMITED

Cambridge England

Published by Woodhead Publishing Limited, Abington Hall, Abington,
Cambridge CB1 6AH, England
www.woodheadpublishing.com

Published in North America by CRC Press LLC, 6000 Broken Sound Parkway, NW,
Suite 300, Boca Raton, FL 33487, USA

First published 2006, Woodhead Publishing Limited and CRC Press LLC
© 2006, Woodhead Publishing Limited
The authors have asserted their moral rights.

This book contains information obtained from authentic and highly regarded sources. Reprinted material is quoted with permission, and sources are indicated. Reasonable efforts have been made to publish reliable data and information, but the authors and the publishers cannot assume responsibility for the validity of all materials. Neither the authors nor the publishers, nor anyone else associated with this publication, shall be liable for any loss, damage or liability directly or indirectly caused or alleged to be caused by this book.

Neither this book nor any part may be reproduced or transmitted in any form or by any means, electronic or mechanical, including photocopying, microfilming and recording, or by any information storage or retrieval system, without permission in writing from Woodhead Publishing Limited.

The consent of Woodhead Publishing Limited does not extend to copying for general distribution, for promotion, for creating new works, or for resale. Specific permission must be obtained in writing from Woodhead Publishing Limited for such copying.

Trademark notice: Product or corporate names may be trademarks or registered trademarks, and are used only for identification and explanation, without intent to infringe.

British Library Cataloguing in Publication Data
A catalogue record for this book is available from the British Library.

Library of Congress Cataloging-in-Publication Data
A catalog record for this book is available from the Library of Congress.

Woodhead Publishing Limited ISBN-13: 978-1-84569-003-8 (book)
Woodhead Publishing Limited ISBN-10: 1-84569-003-6 (book)
Woodhead Publishing Limited ISBN-13: 978-1-84569-173-8 (e-book)
Woodhead Publishing Limited ISBN-10: 1-84569-173-3 (e-book)
CRC Press ISBN-13: 978-0-8493-9159-0
CRC Press ISBN-10: 0-8493-9159-8
CRC Press order number: WP9159

The publishers' policy is to use permanent paper from mills that operate a sustainable forestry policy, and which has been manufactured from pulp which is processed using acid-free and elementary chlorine-free practices. Furthermore, the publishers ensure that the text paper and cover board used have met acceptable environmental accreditation standards.

Project managed by Macfarlane Production Services, Dunstable, Bedfordshire, England
(e-mail: macfarl@aol.com)
Typeset by Godiva Publishing Services Ltd, Coventry, West Midlands, England
Printed by TJ International Limited, Padstow, Cornwall, England

Contents

| | |
|---|----|
| <i>Contributor contact details</i> | xi |
| 1 New brewing technologies: setting the scene | 1 |
| <i>C. W. Bamforth, University of California, USA</i> | |
| 1.1 Introduction | 1 |
| 1.2 The materials used in brewing | 3 |
| 1.3 Brewing issues | 8 |
| 1.4 Sources of further information | 9 |
| 2 Providing cereals for brewing | 10 |
| <i>S. E. Heisel, American Malting Barley Association, USA</i> | |
| 2.1 Introduction | 10 |
| 2.2 Quality evaluations | 11 |
| 2.3 Commercialization of new malting varieties | 20 |
| 2.4 Future trends | 23 |
| 2.5 References | 25 |
| 3 Developments in the supply of adjunct materials for brewing . | 30 |
| <i>D. L. Goode, Kerry Bio-Science, The Netherlands and E. K. Arendt, University College Cork, Ireland</i> | |
| 3.1 Introduction | 30 |
| 3.2 Why use adjuncts? | 30 |
| 3.3 The range of traditional adjuncts | 32 |
| 3.4 Potential new adjunct sources | 43 |
| 3.5 New developments to improve adjunct functionality | 44 |
| 3.6 New beverages based on high adjunct levels | 50 |

| | | |
|----------|---|------------|
| 3.7 | Future trends | 57 |
| 3.8 | Sources of further information | 57 |
| 3.9 | References | 58 |
| 4 | Malt and malt products | 68 |
| | <i>N. Davies, Muntons Plc, UK</i> | |
| 4.1 | Introduction – factors driving change | 68 |
| 4.2 | Malting barley development | 70 |
| 4.3 | Malt processing | 76 |
| 4.4 | Malted ingredients | 92 |
| 4.5 | Sensory evaluation of malt and malted ingredients | 94 |
| 4.6 | Future trends | 97 |
| 4.7 | Sources of further information | 99 |
| 4.8 | References | 100 |
| 5 | The breeding of hop | 102 |
| | <i>J. Henning, Oregon State University, USA</i> | |
| 5.1 | Introduction | 102 |
| 5.2 | Developments in hop breeding | 104 |
| 5.3 | Molecular techniques in hop cultivar development | 106 |
| 5.4 | Conclusion | 118 |
| 5.5 | Sources of further information | 119 |
| 5.6 | References | 119 |
| 6 | The processing of hops | 123 |
| | <i>C. Schönberger, Joh. Barth & Sohn GmbH & Co. KG, Germany</i> | |
| 6.1 | Introduction – the processing of hops | 123 |
| 6.2 | Hop pellets | 125 |
| 6.3 | Hop extracts | 126 |
| 6.4 | Isomerised hop products | 132 |
| 6.5 | Other hop products | 138 |
| 6.6 | Hop, hop products and relevant beer analyses | 142 |
| 6.7 | Future trends | 145 |
| 6.8 | Sources of further information | 145 |
| 6.9 | References | 146 |
| 7 | Yeast genetics in brewing: new insights and opportunities | 149 |
| | <i>D. E. Quain, red.ts Ltd, UK</i> | |
| 7.1 | Introduction | 149 |
| 7.2 | Fundamentals | 151 |
| 7.3 | The yeast genome | 151 |
| 7.4 | The rise and fall of genetic modification | 154 |
| 7.5 | Genetic instability – problem or opportunity? | 157 |
| 7.6 | Genetic differences between yeast species | 159 |
| 7.7 | Future trends | 163 |

| | | |
|-----------|--|------------|
| 7.8 | Sources of further information | 163 |
| 7.9 | Acknowledgements | 163 |
| 7.10 | References | 163 |
| 8 | Yeast supply and propagation in brewing | 167 |
| | <i>D. Quain, Coors Brewers Ltd, UK</i> | |
| 8.1 | Introduction | 167 |
| 8.2 | Storage and supply of yeast cultures | 168 |
| 8.3 | Propagation | 171 |
| 8.4 | Active dried yeasts | 178 |
| 8.5 | Future trends | 179 |
| 8.6 | Sources of further information | 180 |
| 8.7 | Acknowledgements | 180 |
| 8.8 | References | 180 |
| 9 | Water in brewing | 183 |
| | <i>M. Eumann, EUWA Water Treatment Plants, Germany</i> | |
| 9.1 | Introduction | 183 |
| 9.2 | Water for use in breweries | 184 |
| 9.3 | Treatment technologies | 190 |
| 9.4 | Recycling | 205 |
| 9.5 | Future trends | 206 |
| 9.6 | Sources of further information | 206 |
| 9.7 | Bibliography | 207 |
| 10 | The brewhouse | 208 |
| | <i>J. M. H. Andrews, Briggs of Burton Plc, UK</i> | |
| 10.1 | Introduction | 208 |
| 10.2 | Milling | 209 |
| 10.3 | Mashing devices | 212 |
| 10.4 | Mash conversion vessels | 213 |
| 10.5 | Mash separation | 215 |
| 10.6 | Wort boiling | 221 |
| 10.7 | Energy recovery systems | 224 |
| 10.8 | Hot wort clarification | 225 |
| 10.9 | Wort cooling | 226 |
| 10.10 | Cold wort clarification | 226 |
| 10.11 | Cold wort oxygenation | 227 |
| 10.12 | References | 227 |
| 11 | Fermentation of beer | 228 |
| | <i>C. Boulton, Coors Brewers Ltd, UK</i> | |
| 11.1 | Introduction | 228 |
| 11.2 | Current developments | 229 |
| 11.3 | Fermentation management | 230 |

| | | |
|-----------|---|------------|
| 11.4 | Yeast distribution in the fermenter | 239 |
| 11.5 | CO ₂ collection | 241 |
| 11.6 | Modelling fermentation | 242 |
| 11.7 | Continuous fermentation | 244 |
| 11.8 | High gravity brewing | 246 |
| 11.9 | Yeast physiology and fermentation performance | 247 |
| 11.10 | Future trends | 249 |
| 11.11 | Sources of further information | 250 |
| 11.12 | References | 251 |
| 12 | Accelerated processing of beer | 254 |
| | <i>I. Virkajärvi, VTT Technology, Finland</i> | |
| 12.1 | Introduction | 254 |
| 12.2 | Diacetyl reduction | 255 |
| 12.3 | New techniques | 256 |
| 12.4 | Warm stabilization | 265 |
| 12.5 | Cold stabilization | 266 |
| 12.6 | Two maturation philosophies | 268 |
| 12.7 | Sources of further information | 269 |
| 12.8 | References | 270 |
| 13 | Filtration and stabilisation of beer | 275 |
| | <i>G. Freeman, Brewing Research International, UK</i> | |
| 13.1 | Introduction | 275 |
| 13.2 | Current filtration practice | 275 |
| 13.3 | Possibilities in recycling of kieselguhr | 278 |
| 13.4 | Crossflow filtration | 279 |
| 13.5 | Single-pass membrane filtration | 282 |
| 13.6 | Novel filter aids | 284 |
| 13.7 | Current stabilisation processes | 285 |
| 13.8 | Novel stabilising systems | 287 |
| 13.9 | The effect of modern processing technologies on stability ... | 288 |
| 13.10 | Pasteurisation | 289 |
| 13.11 | Future trends | 290 |
| 13.12 | Acknowledgement | 291 |
| 13.13 | References | 291 |
| 14 | Packaging of beer | 293 |
| | <i>J. Browne, Technical Management Support Ltd, UK</i> | |
| 14.1 | Introduction | 293 |
| 14.2 | Trends in packaging formats | 293 |
| 14.3 | Developments in canning and bottling | 298 |
| 14.4 | Developments in handling kegs | 303 |
| 14.5 | Future trends | 305 |
| 14.6 | Sources of further information | 307 |

| | | |
|-----------|---|------------|
| 14.7 | Acknowledgements | 307 |
| 14.8 | References | 307 |
| 15 | Modern brewery sanitation | 308 |
| | <i>D. Loeffler, Loeffler Chemical Corporation, USA</i> | |
| 15.1 | Introduction | 308 |
| 15.2 | Sanitation terminology | 309 |
| 15.3 | Theoretical aspects of cleaning | 311 |
| 15.4 | The use of water in cleaning | 313 |
| 15.5 | Detergents/cleaners | 314 |
| 15.6 | Mechanical aspects of cleaning | 319 |
| 15.7 | Temperature | 326 |
| 15.8 | Time | 327 |
| 15.9 | Sanitizers/disinfectants | 328 |
| 15.10 | Future trends | 333 |
| 15.11 | Sources of further information | 334 |
| 16 | Waste handling in the brewing industry | 335 |
| | <i>R. Reed, Black & Veatch, UK</i> | |
| 16.1 | Introduction | 335 |
| 16.2 | Assessment of the character and strength of wastewaters | 336 |
| 16.3 | Sources and nature of wastewater | 337 |
| 16.4 | Extent of treatment | 341 |
| 16.5 | Outline of processes and technology | 342 |
| 16.6 | Treatment processes | 346 |
| 16.7 | Summary of the new legislation | 353 |
| 16.8 | Management approaches to water and waste minimisation ... | 355 |
| 16.9 | Future trends | 356 |
| 16.10 | References | 357 |
| 17 | Quality assurance in brewing | 358 |
| | <i>G. Jackson, Brewing Research International, UK</i> | |
| 17.1 | Introduction | 358 |
| 17.2 | Hazard Analysis Critical Control Points (HACCP) | 358 |
| 17.3 | Quality management systems and ISO9001:2000 | 363 |
| 17.4 | Integrated management systems – the BRC Global Standard – Food | 366 |
| 17.5 | Feed Materials Assurance Scheme (FEMAS) | 368 |
| 17.6 | Future trends | 371 |
| 17.7 | References | 371 |
| 18 | Brewing control systems: chemical analytes | 372 |
| | <i>K. J. Siebert, Cornell University, USA</i> | |
| 18.1 | Introduction | 372 |
| 18.2 | Brewing analytical methods | 373 |
| 18.3 | Detection, separation and measurement techniques | 373 |

| | | |
|--------------|--|------------|
| 18.4 | Combining different techniques | 379 |
| 18.5 | What and why do we measure? | 382 |
| 18.6 | Where and how do we measure? | 384 |
| 18.7 | Impact of brewery operation scale | 385 |
| 18.8 | Changes over time | 385 |
| 18.9 | Traditional, emerging and future methods | 386 |
| 18.10 | New technologies | 387 |
| 18.11 | References | 387 |
| 19 | Brewing control systems: microbiological analysis | 391 |
| | <i>E. Storgårds, A. Haikara and R. Juvonen, VTT Technical Research Centre of Finland</i> | |
| 19.1 | Introduction | 391 |
| 19.2 | Classical microbiological methods | 397 |
| 19.3 | Optical techniques | 400 |
| 19.4 | Molecular methods for detection and identification | 403 |
| 19.5 | Indirect methods | 407 |
| 19.6 | Evaluation of yeast quality and quantity | 409 |
| 19.7 | Future trends | 413 |
| 19.8 | Sources of further information | 415 |
| 19.9 | References | 416 |
| 20 | Brewing control systems: sensory evaluation | 427 |
| | <i>W. J. Simpson, FlavorActiV Ltd, UK</i> | |
| 20.1 | Introduction | 427 |
| 20.2 | Brands | 427 |
| 20.3 | Tasters | 431 |
| 20.4 | Assessment methods | 442 |
| 20.5 | Assessment facilities | 448 |
| 20.6 | Data analysis tools | 449 |
| 20.7 | Data reporting and distribution tools | 450 |
| 20.8 | Future trends | 450 |
| 20.9 | Sources of further information | 451 |
| 20.10 | Acknowledgements | 453 |
| 20.11 | References | 453 |
| 21 | The future of brewing | 461 |
| | <i>C. W. Bamforth, University of California, USA</i> | |
| 21.1 | Choice | 461 |
| 21.2 | Impact of choice | 462 |
| 21.3 | Technical need drives research | 462 |
| 21.4 | Global influences | 465 |
| 21.5 | The nature of the brewing process in 2050 | 467 |
| <i>Index</i> | | 468 |

Contributor contact details

(* = main contact)

Chapters 1 and 21

Dr C. W. Bamforth
Chair and Professor
Department of Food Science and
Technology
University of California
Davis, CA 95616
USA
E-mail: cwbamforth@ucdavis.edu

Chapter 2

Mr S. E. Heisel
American Malting Barley Association
740 N Plankinton Ave
Suite 830
Milwaukee, WI 53203-2403
USA

Chapter 3

Dr D. L. Goode*
Kerry Bio-Science BV
Veluwezoom 62
1327 AH Almere
The Netherlands
E-mail:
declan.goode@kerrybioscience.com

Dr E. K. Arendt
Department of Food and Nutritional
Sciences
University College Cork
Cork
Ireland
E-mail: e.arendt@ucc.ie

Chapter 4

Dr N. Davies
Technical Director
Muntons Plc
Cedars Maltings

Stowmarket
Suffolk IP14 2AG
UK
E-mail: nigel.davies@muntons.com

Chapter 5

Dr J. Henning
Research Hop Geneticist (plants)
USDA-ARS
3450 SW Campus Way
Corvallis, OR 97330
USA
E-mail: john.Henning@orst.edu

Chapter 6

Dr C. Schönberger
Joh. Barth & Sohn GmbH & Co. KG
Freiligrathstrasse 7/9
90482 Nuernberg
Germany
E-mail:
Christina.Schoenberger@johbarth.de

Chapters 7 and 8

Dr D. E. Quain
red.ts Ltd
9 Wheatfield Court
Willington
Derbyshire DE65 6PT
UK
E-mail: david.quain@red.ts.com

Chapter 9

M. Eumann
EUWA H.H. Eumann GmbH
Daimlerstrasse 2-10
D-71116 Gärtringen
Germany
E-mail: info@euwa.com

Chapter 10

Dr J. M. H. Andrews
Briggs of Burton Plc
Briggs House
Derby Street
Burton-on-Trent
Staffordshire DE14 2LH
UK
E-mail: jmha@briggsplc.co.uk

Chapter 11

Dr C. Boulton
Coors Brewers Ltd
Technical Centre
PO Box 12
Cross Street
Burton-on-Trent
Staffordshire DE14 1XH
UK
E-mail:
chris.boulton@coorsbrewers.com

Chapter 12

Dr I. Virkajärvi
Tuohimäki 3 B6
FIN-02130 Espoo
Finland
E-mail: ikka.virkajarvi@vtt.fi

Chapter 13

Mr G. Freeman
Brewing Research International
Coopers Hill Road
Nutfield
Redhill
Surrey RH1 4HY
UK

E-mail:
g.freeman@brewingresearch.co.uk

Chapter 14

Mr J. Browne
Technical Management Support Ltd
Claim Farm House
Manley Road
Kingswood
Frodsham
Cheshire WA6 6HT
UK

E-mail:
jeremy.browne@btinternet.com

Chapter 15

Mr D. Loeffler
Technical Director
Loeffler Chemical Corporation
5700 Bucknell Drive
Atlanta, GA 30336
USA

E-mail:
loeffler.dirk@loefflerchemical.com

Chapter 16

Dr R. Reed
Brackenbury House
Lodge Lane

Salfords
Redhill
Surrey RH1 5DH
UK
E-mail: reedr@bv.com

Chapter 17

Mr G. Jackson
Brewing Research International
Coopers Hill Road
Nutfield
Redhill
Surrey RH1 4HY
UK

E-mail:
g.jackson@brewingresearch.co.uk

Chapter 18

Dr K. J. Siebert
Food Science and Technology
Department
Cornell University
Geneva, NY 14456
USA

E-mail: kjs3@cornell.edu

Chapter 19

Dr E. Storgårds*, A. Haikara and
R. Juvonen
VTT Technical Research Centre of
Finland
PO Box 1000
FI-02044 VTT
Finland

E-mail: erna.storgards@vtt.fi

Chapter 20

Dr W. J. Simpson
FlavorActiV Limited
Sanderum House
Oakley Road
Chinnor
Oxfordshire OX39 4TW
UK

E-mail: bill.simpson@flavoractiv.com

Kerry Bio-Science, your brewing partner

Kerry Bio-Science is one of the market leaders in innovation and the application of bio- ingredients for brewing, distilling and beverage industries. Our range of brewing process aids and ingredients compliment the natural brewing process to produce beer of consistent quality. Our experienced team of experts help customers to create quality products that meet the rapidly changing demands of today's consumer market.

Kerry Bio-Science offers ingredients that can help you in every stage of the brewing process:

ENZYMES

| | |
|-------------------|--|
| Type of enzymes | : amylase, protease; glucanase for all brewing applications giving improved brewhouse performance (filtration and extract yield) |
| Functionality | |
| Range of products | : Bioglucanase, Bioamylase, Bioprotease, Promalt and Hitampase |

FINING AGENTS

| | |
|-------------------|-----------------------------------|
| Functionality | : for wort and beer clarification |
| Range of products | : Whirlfloc and Bioline |

FOAM STABILISERS

| | |
|-------------------|----------------------|
| Functionality | : foam stabilisation |
| Range of products | : Biofoam |

FLAVOUR STABILITY

| | |
|-------------------|-------------------------|
| Type of products | : anti-oxidant products |
| Range of products | : Blox |

FERMENTATION AIDS

| | |
|-------------------|-------------------------------------|
| Functionality | : optimize fermentation performance |
| Range of products | : Yeastex and Fermcap |



KERRY

Kerry Bio-Science

Brewing Technology Centre
Kerry Bio-Science Ireland
Kinagleary, Carrigaline
Co. Cork, Ireland
T +353 214 376400 / F +353 214 376480



1

New brewing technologies: setting the scene

C. W. Bamforth, University of California, USA

1.1 Introduction

The aim of this book is not primarily to tackle the science underpinning malting and brewing. Rather, the focus is on practical issues. In this chapter I will set the scene with some underpinning information, but those seeking the necessary basic scientific descriptions of everything from barley to beer should consult a text such as Bamforth (2006) or, at a more advanced level, Briggs *et al.* (2005). For immediate purposes Table 1.1 offers basic summaries of the processes involved in malting and brewing and the importance of each unit operation.

Table 1.1 The essentials of malting and brewing

| Process stage | Description |
|---|---|
| <i>Selection of barley</i> ¹ | Malting barleys (moisture content <12%) of relatively low total N content (e.g. less than 1.7% N), with high viability and with endosperm of mealy texture that hydrates readily and possesses cell walls that are readily degraded |
| <i>Storage of barley</i> | Sometimes important to allow barley to free up from innate dormancy |
| <i>Malting: steeping</i> | Staged addition of water at 14–18°C separated by air rests to raise moisture content to 43–46%. Allows the embryo to start synthesising hormones (notably gibberellins); allows the aleurone to become receptive to the hormone action that will trigger enzyme synthesis; allows the starchy endosperm to become receptive to digestion ('modification') |

2 Brewing

Table 1.1 (continued)

| Process stage | Description |
|---|--|
| <i>Malting: germination</i> | Controlled germination for 3–6 days at 16–20°C to degrade the endosperm cell walls and much of the protein. Enzymes synthesised by the aleurone migrate from proximal to distal end, digesting cell walls (β -glucanase family of enzymes, pentosanases) and protein (proteinase family). Causes softening of grain. Amylases also developed or activated, but limited in action on starch granules |
| <i>Malting: kilning</i> | Drying of malt at successively increasing temperatures (mainstream malts max. 105°C) to dry the grain (target <6% H ₂ O), whilst retaining much enzymic activity and developing colour and flavour through Maillard interactions between sugars and amino acids produced during modification |
| Malt: storage | 2–4 weeks storage to avoid wort separation problems in the brewery |
| Milling and mashing | Generation of particles accessible to mashing water, mashing often starting at say 50°C (20 minutes) to allow remaining action of thermolabile β -glucanase, then passing through 65°C (e.g. for 1 h) for starch gelatinisation and action of amylase complex. Wort produced by conventional mashing has 20–25% starch left behind as non-fermentable digestion products called dextrans. Wort separated from grains |
| Boiling | Wort boiled with hops or hop preparations, typically for 1 hour. Isomerisation of bitter acids (main part of resin fraction) to increase their solubility and bitterness. Volatilisation of aroma components, including hop oils unless hops added late in the boil. Clarification stage follows to remove ‘hot break’ and residual hop material. Wort cooled and air or oxygen added |
| Fermentation | Ales typically fermented warmer (15–25°C) and therefore faster than lagers (6–15°C). Time range 3–14 days. Fermentation at a targeted rate of specific gravity drop and to a target ‘attenuation’. Also diacetyl and pentanedione, which afford butterscotch/honey aromas, must be removed by prolonged contact of yeast with ‘green beer’ |
| Maturation, stabilisation and packaging | Minimum regime is –1°C, usually for 2–3 days. Some hold longer. Insolubilisation and settling of proteins and polyphenols. Filtration (kieselguhr or perlite-based). Removal of residual haze precursors by polyvinylpyrrolidone (polyphenols) and/or silica hydrogels or tannic acid or papain (proteins). Removal of any microbial contamination by pasteurisation or filtration. Adjustment of CO ₂ content, then fill vessels |

¹ *Process stages in italics occur in the maltings, prior to the brewery.*

Source: Bamforth, C.W. (2004) Opportunities for newer technologies in the oldest biotechnology, brewing. *Applied Biotechnology, Food Science and Policy*, 1, 213–222.

1.2 The materials used in brewing

The larger proportion of the world's beers is produced from *malted barley*. It has been this way for perhaps 8000 years. Barley has been retained as the primary cereal of choice, not least because it retains its husk on threshing and this traditionally form the filter bed through which wort is collected in the brew house. As we shall see in Chapter 10, the mash filter does not depend on the husk in this way, which might in future open up possibilities for huskless barleys for brewing. However, these handle in the maltings much as does *wheat*, with a tendency towards stickiness. Wheat is the second most employed cereal in brewing, notably for the production of weissbiers and weizenbiers in Germany.

Barley is malted prior to use, in order that the enzymes that degrade starch to fermentable sugars are synthesised. Additionally there is a synthesis of the enzymes that degrade the cell walls and much of the protein in the starchy endosperm, thereby softening the grain and making it more millable. There is an unavoidable development of embryo tissue (rootlets and acrospire), and the maltster seeks to balance the extent of this with the need for adequate 'modification' of the endosperm.

Although there are beers that are produced only from malted barley e.g. those produced under the terms of the five-centuries-old Bavarian 'purity' law, the *Reinheitsgebot*, many brewers employ various adjuncts for reasons of quality (different colours, better foams, interesting flavours) or cost. In fact there may not be quite the cost savings anticipated from use of an ostensibly cheaper starch source, because the brewer may have to employ more expensive processing procedures if problems are to be avoided. The major cost components in brewing are illustrated in Fig. 1.1. It will be apparent that the costs of packaging, processing, taxation and marketing vastly exceed those of raw materials. It is only when there are political reasons or taxation reasons – e.g. Happoshu, see Chapter 3) – that there are major justifications to be made on a cost basis for the use of adjuncts.

Of particularly low relative cost to the brewer are the *hops*, yet these make a huge contribution to product quality and stability: apart from the bitterness from the resins and aroma from the oils, the resins also afford anti-microbial properties and foam stabilisation but also comprise precursors of staling and of the skunky aromas that develop in beer exposed to light. Despite the low cost of hops relative to that of the beer, there is a plethora of hop preparations available to the brewer, including those that specifically protect against this light sensitivity (Chapter 6).

The bulk of most beers comprises *water*, hence the scrutiny which brewers devote to this product. Vastly more water is needed to make a pint of beer than actually finds its way into the beer, perhaps five times more for a well-run brewery and 20 times more for a badly operated facility.

Most brewers maintain their own *yeast* strains. As the alcohol concentration of most beers does not become too high during fermentation, the yeast that multiplies in fermentation remains healthy and suitable for re-pitching into



Fig. 1.1 A generalised breakdown of major costs associated with the brewing of beer.

subsequent fermentations. Some smaller brewers have done this in seeming perpetuity; however, the latter-day 'gold standard' is to re-pitch with newly propagated yeast every fourth or fifth fermentation. There is some interest in the use of dried yeast preparations of the type used *inter alia* in baking and wine making (Chapter 8).

As they will not be addressed elsewhere in this book, let us now consider miscellaneous process ingredients that may be employed in brewing. Not included in the discussion are the salts that may be added to adjust the ionic balance of the grist, e.g. calcium salts to increase the hardness ('Burtonisation'), and the zinc chloride or zinc sulphate that many brewers add to encourage yeast activity.

1.2.1 Exogenous enzymes

Various 'brewing enzymes' have been marketed over the years with the intent of satisfying various demands of the brewer (Table 1.2). The majority of these enzymes are not pure, insofar as they contain a range of enzymic activities. During the growth of the organism, different enzymes will be successively released into the medium in the order in which they are needed. So the precise balance of enzymes that is packaged and made available to the brewer will very much depend on where the 'cut' was taken. Nowadays most of the brewing enzymes are made using organisms that are genetically modified so as to optimise the level and type of enzymes in the broth.

1.2.2 Isinglass

One of the great beer genres (*viz.* the English cask ale) emerged on the backbone of a 'natural' clarification process rooted in a protein preparation called isinglass. Isinglass is a very pure form of collagen obtained from the dried swim bladders (some call them 'maws') of certain warm-water fish, amongst them the catfish, jewfish, threadfish and croaker. These fish are primarily caught for food use and the functional property of the maw represents added value.

Table 1.2 Exogenous enzymes used in brewing

| Enzyme | Stage of use | Function |
|---------------------------------|-------------------------|---|
| β -Glucanase ¹ | Mashing | Eliminate glucans that cause wort separation, filtration and clarity problems |
| Pentosanase ¹ | Mashing | Support glucanases in digestion of cell wall polymers from barley and wheat |
| Proteinase ¹ | Mashing | Ensure generation of sufficient yeast-assimilable amino acids |
| α -Amylase ¹ | Mashing | Starch digestion |
| Glucoamylase | Mashing or fermentation | Enhance starch digestion, to the extent of allowing conversion to totally fermentable sugars of value in the production of light and low carbohydrate beers |
| Pullulanase | Mashing | Promote digestion of branched-chain dextrins |
| Acetolactate decarboxylase | Fermentation | Accelerate elimination of vicinal diketones in beer maturation |
| Papain | Beer in storage | Eliminate haze-forming polypeptides |
| Prolyl endopeptidase | Beer in storage | Selectively remove haze-forming polypeptides; of potential value in producing beer for coeliacs |

¹ Of especial significance if malt replaced by high levels of grain adjunct, e.g. unmalted barley.

The bladders are removed, washed and dried. At the smallest scale in a fishing village the maws are sun-dried, but modern fish processing plants use commercial dryers. Dried maws are ground up, washed and sterilised before being 'cut' by weak acids such as sulphurous acid to disrupt the structure of the collagen molecules so as to generate the correct balance and orientation of positively and negatively charged sites that are responsible for its functionality.

Worldwide, few brewers use isinglass, despite the fact that it is very effective in settling the contents of conditioning tanks, thereby minimising the loading of solids onto the filter without the expense of a centrifuge.

Various other fining agents are sometimes used in the brewing industry. These may be classified as kettle (or copper) finings, which are added during the boil, and auxiliary finings, which are used alongside isinglass to aid clarification of beer. The best-known kettle fining agent is the negatively charged carrageenan (Irish Moss), derived from seaweed, and the most frequently used auxiliary finings are silicates and alginates, the former derived from sand and the latter from seaweed. Again, both are negatively charged and complement the action of the positively charged isinglass.

1.2.3 Filter aids

There are fundamentally two types of filter aid: kieselguhr or perlite. Kieselguhr, or 'diatomaceous earth', comprises silica-based shells of ancient unicellular aquatic microscopic plants called diatoms. Its heat resistance means that it can

be used as an insulator, but its abrasiveness means that it has also formed a component of toothpaste and metal polishes. Apart from being widely used as a filter aid to clarify syrups as well as alcoholic beverages, it is used as a filling material in paper, paints, ceramics, soap and detergents. Alfred Nobel found that it is a great absorbant of nitro-glycerine in the manufacture of dynamite.

Huge beds of kieselguhr, between 40 and 50 feet (12–15 metres) deep, are found in Virginia, but also in parts of Germany and in Aberdeenshire in Scotland. The microscopic appearance from different localities differs considerably. The deposits contain varying amounts of organic matter together with sand, clay, and iron oxide, and the raw material is first incinerated (calcined) to destroy organic matter. The successive process stages in rendering bags of kieselguhr in the form needed by the brewer are mining, crushing, drying, calcining, cooling, air classification and packaging.

The interests of purveyors of perlite are looked after by the Perlite Institute (go to <http://www.perlite.org/>). Perlite is a naturally occurring siliceous rock that, when heated, expands from four to 20 times its original volume. When heated to above 871°C, it pops like popcorn to produce many small bubbles, so perlite is very light and white.

There are many uses for perlite. Its insulating properties and lightness render it valuable as an insulator in masonry and cryogenic vessels. It is used as an aggregate in cement and plasters and for under-floor insulation, chimney linings, paint texturing, gypsum boards, ceiling tiles, and roof insulation boards. Perlite is used as a component of soil-less growing mixes, allowing aeration and moisture retention. It is also used as a carrier for fertiliser, herbicides and pesticides and for pelletising seed.

Apart from clarifying beer, perlite is also used for cleaning up pharmaceuticals, chemicals and water. Like kieselguhr, it can also be used as an abrasive.

1.2.4 Stabilisers

Silica hydrogels and xerogels have their origins in a pure form of sand. This is first converted into a soluble form by the action of alkali. Thereafter there is a controlled aggregation of sodium silicate from a sol form under acidic conditions. The washed aggregated particles are processed by techniques including micronisation, drying, milling and classifying to yield the desired balance of particle sizes and pore sizes, the range of which is taken advantage of by brewers (and others) to remove colloidal particles of various types from their products.

Polyvinylpyrrolidone (PVPP) is produced by a technique called 'popcorn polymerisation'. The monomer vinylpyrrolidone is heated with strong caustic and then cooled, in which phase the polymerisation takes place. Subsequently there are slurring, filtering, hydrolysis (using phosphoric acid), washing, re-slurring, and drying stages, in which any residual monomer and water are removed.

Alternative stabilisers are now in the market, including a PVP–silica gel composite.

1.2.5 Foam stabiliser

Propylene glycol alginate (PGA) is rather more relevant as a foam *protectant* than as a foam stabiliser. It helps prevent lipid and detergent damage at point of sale, without offering absolute protection. Alginates are polysaccharides extracted by alkali from brown seaweed. They have been known since their discovery by an English chemist in 1881. The seaweed is harvested on the coasts of North America, Scotland, Ireland, Norway, France, Japan, China, Korea, Chile, South Africa and Australia. Of these China easily generates the most, though the UK and the USA are the biggest processors, with one company dominating, at 70% of the market.

Insoluble alginate from the raw plant is solubilised by ion exchange, then filtered. The alginic acid obtained is partially dehydrated and then esterified by reaction with gaseous propylene oxide under pressure at 45–60°C. After washing with alcohol, the product is dried and milled.

Table 1.3 Issues of quality in brewing

| Issue | Description |
|--------------------|---|
| Foam | Most beers are expected to have an appealing, stable foam. However, increasing tendency for consumers to drink directly from the container. Impacting factors include amphipathic polypeptide survival from the grist, presence of bitter acids (reduced side-chain acids give more stable foams), levels of CO ₂ and N ₂ , nucleation devices, absence of lipids and detergents. |
| Clarity | Most beers are bright (free from haze or sediment) – but not all. Insoluble materials can have diverse origins, including proteins cross-linking with oxidised polyphenols, residual grain polysaccharides, oxalate, micro-organisms |
| Colour | Derives from Maillard reaction in malt kilning and adjunct roasting and from polyphenol oxidation in the brew house |
| Absence of gushing | Principal cause is <i>Fusarium</i> infection of grain, with release of hydrophobin. But can also be cause by oxidised resins in hop bitterness preparations and insolubles in beer, e.g. oxalate |
| Package | Major determinant of impression customer gets of product – e.g. absence of scuffing on bottles, making returnable glass less appealing than one-trip glass |
| Flavour | Impacted by all the above. Pronounced marketing pressure to have beer in any coloured glass other than brown, but flint and green glass permit light passage that causes skunking of beer. Flavour impacted by all raw materials, demanding control of material selection and processing. Flavour instability the major unsolved problem facing brewers, in particular development of papery flavour. Stringent control of oxygen a major factor, but not alone. Cold shipping and stockholding of beer expensive but effective |

Note: enhanced analytical capabilities (e.g. sensors) will permit increased control, with impacts on cost and quality.

sample content of Brewing: New Technologies (Woodhead Publishing Series in Food Science, Technology and Nutrition)

- [Lacanian Ethics and the Assumption of Subjectivity pdf, azw \(kindle\)](#)
- [read online Cocaine Wars](#)
- **[download The Art of The Last of Us](#)**
- [read Ivar's Seafood Cookbook: The O-fish-al Guide to Cooking the Northwest Catch](#)

- <http://www.gateaerospaceforum.com/?library/Beer-O-Clock--An-Insider-s-Guide-to-History--Craft--and-Culture.pdf>
- <http://aircon.servicessingaporecompany.com/?lib/The-Souls-of-Black-Folk--Oxford-World-s-Classics-.pdf>
- <http://honareavalmusic.com/?books/The-Art-of-The-Last-of-Us.pdf>
- <http://reseauplatoparis.com/library/The-Original-Curse--Did-the-Cubs-Throw-the-1918-World-Series-to-Babe-Ruth-s-Red-Sox-and-Incite-the-Black-Sox-Sc>